

University of Groningen

Antibiotic resistance, stewardship, and consumption

Pouwels, Koen B.; Chatterjee, Anuja; Cooper, Ben S.; Robotham, Julie V.

Published in:
The Lancet Planetary Health

DOI:
[10.1016/S2542-5196\(18\)30283-3](https://doi.org/10.1016/S2542-5196(18)30283-3)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2019

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Pouwels, K. B., Chatterjee, A., Cooper, B. S., & Robotham, J. V. (2019). Antibiotic resistance, stewardship, and consumption. *The Lancet Planetary Health*, 3(2), E66. [https://doi.org/10.1016/S2542-5196\(18\)30283-3](https://doi.org/10.1016/S2542-5196(18)30283-3)

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Antibiotic resistance, stewardship, and consumption

Peter Collignon and colleagues¹ aim to improve understanding of global antibiotic resistance drivers by quantifying associations between antibiotic resistance and potential contributing factors. A recent systematic review including 565 studies primarily reporting patient-level data found that antibiotic exposure, underlying disease, and invasive procedures had the greatest evidence of being antibiotic resistance drivers.² Given the small evidence base for the role of community factors,² we fully concur with Collignon and colleagues about the importance of considering such factors in the spread of antibiotic resistance.¹ However, as the authors recognise, correlation does not imply causation—indeed, national meticillin-resistant *Staphylococcus aureus* rates strongly correlate with unfair play of national football teams.³ Conversely, it would be a mistake to conclude that weak (or even inverse) correlation between antibiotic use and antibiotic resistance at the national level implies an absence of a strong causal relationship (appendix).

There are several limitations to using current global data. First, important between-country differences in the quality and representativeness of data are likely to exist. For example, blood cultures are less frequently taken in resource-poor settings than in resource-rich settings, and often only when patients do not respond to empirical therapy, leading to inflated estimates of resistance.⁴ Moreover, in lower income countries, a substantial proportion of antibiotics used might come from informal providers and not appear in consumption data. Both factors would be expected to weaken associations between reported antibiotic use and resistance, and could lead to positive associations between

governance and infrastructure and antibiotic resistance, even if these factors have no effect on antibiotic resistance.

Additionally, comparisons based on national-level data can be problematic, particularly when many countries have a sharp divide between subpopulations that overuse antibiotics and those that are without access. Patients lacking access to antibiotics are also less likely to have access to health-care facilities where blood samples are taken. Considering antibiotic consumption only on a national per capita basis could therefore considerably underestimate antibiotic consumption in subpopulations where resistance is being measured. Using a multivariable model will unfortunately not overcome such biases.

Finally, the analysis simplifies antibiotic use such that the matter of which antibiotic is consumed is not considered, which will have importance in terms of resistance development.⁵ Combined antibiotic consumption measures might therefore not be optimal for determining the contribution of antibiotic consumption to resistance.

While we welcome the central conclusion of the study about the need for efforts to reduce the spread of antibiotic resistance in the community, we are concerned that fundamental limitations of the data could have led to an undervaluing of the role of antibiotic stewardship.

We declare no competing interests.

Copyright © 2019 The Author(s). Published by Elsevier Ltd. This is an Open Access article under the CC BY 4.0 license.

***Koen B Pouwels, Anuja Chatterjee, Ben S Cooper, Julie V Robotham
koen.pouwels@phe.gov.uk**

Modelling and Economics Unit, National Infection Service, Public Health England, London NW9 5EQ, UK (KBP, JVR); Unit of Global Health, Department of Health Sciences, University Medical Centre Groningen, University of Groningen, Groningen, Netherlands (KBP); National Institute for Health Research, Health Protection Research Unit in Healthcare Associated Infection and Antimicrobial Resistance, Imperial College London, London, UK (AC, JVR); Centre for Tropical Medicine and Global

Health, Nuffield Department of Medicine, University of Oxford, Oxford, UK (BSC); and Mahidol Oxford Tropical Medicine Research Unit, Faculty of Tropical Medicine, Mahidol University, Bangkok, Thailand (BSC)

- 1 Collignon P, Beggs JJ, Walsh TR, Gandra S, Laxminarayan R. Anthropological and socioeconomic factors contributing to global antimicrobial resistance: a univariate and multivariable analysis. *Lancet Planet Health* 2018; **2**: e398–405.
- 2 Chatterjee A, Modarai M, Naylor NR, et al. Quantifying drivers of antibiotic resistance in humans: a systematic review. *Lancet Infect Dis* 2018; **18**: e368–78.
- 3 Meyer E, Gastmeier P, Schwab F. National MRSA rates run along with fair play of national football teams: a cross-sectional data analysis of the European Football Championship, 2008. *Infection* 2013; **41**: 215–18.
- 4 De Kraker ME, Stewardson AJ, Harbarth S. Will 10 million people die a year due to antimicrobial resistance by 2050? *PLoS Med* 2016; **13**: e1002184.
- 5 Pouwels KB, Freeman R, Muller-Pebody B, et al. Association between use of different antibiotics and trimethoprim resistance: going beyond the obvious crude association. *J Antimicrob Chemother* 2018; **73**: 1700–07.



See Online for appendix